

# TRYPTONE WATER

A differential medium for the differentiation of bacteria on the basis of their ability to produce indole from tryptophan.

<b>Dehydrated media</b>	
Code number:	500 g: TRW20500, 5 kg: TRW25000
Colour:	Yellowish
Appearance:	Homogeneous hygroscopic powder
pH before autoclaving (25 °C):	7,3 – 7,7

**Direction:** Suspend **15 g** in one litre of distilled water and heat gently to dissolve the medium completely. Dispense into test tubes and sterilise by autoclaving at 121 °C for 15 minutes.

<b>Prepared media</b>	
Bottled media:	100 ml: TRW30100, 500 ml: TRW30500
Tubed media:	100 x 12 mm: TRW40003 (3 ml)
Colour:	Yellowish
pH (25 °C):	7,4 – 7,6

**Direction:** Dispense the bottled media aseptically into sterile test tubes. Media in tubes are ready to use.

## FORMULA in g/l

Tryptone	10
Sodium chloride	5

**Note:** The typical formula can be adjusted to obtain optimal performance.

**Storage conditions:** Store the dehydrated media tightly closed in a dry place at room temperature. Store the bottled and tubed media protected from light at room temperature. Use before the expiry date on the label.

### Quality control:

<b>Test strains</b>	Incubation temp: 37 °C	<b>Growth</b>	Incubation time: 24 h
<i>Escherichia coli</i>	ATCC 25922	Indole positive	
<i>Enterobacter aerogenes</i>	ATCC 13048	Indole negative	

**References:** Farmer (1985) J. Clin. Microbiol. 21: 46.

**In vitro diagnostic – for professional use only!**